

January 23, 2023

Honorable Kathy Hochul
Governor
New York State Capitol
Albany, NY 12224

Honorable Pete Harckham
Chair, Senate Environmental Conservation Committee
New York State Senate
Albany, NY 12247

Honorable Andrea Stewart-Cousins
President Pro Tempore and Majority Leader
New York State Senate
Albany, NY 12247

Honorable Deborah Glick
Chair, Assembly Environmental Conservation
Committee
New York State Assembly
Albany, NY 12248

Honorable Carl Heastie
Speaker
New York State Assembly
Albany, NY 12248

Honorable Brad Hoylman-Sigal
Chair, Senate Judiciary Committee
New York State Senate
Albany, NY 12247

Dear Governor Hochul, President Pro Tempore and Majority Leader Stewart-Cousins, Speaker Heastie, and Chairs Harckham, Glick, and Hoylman:

The undersigned environmental health scientists and health professionals urge you to eliminate harmful and unnecessary uses of neonicotinoid pesticides (neonics) in New York State—specifically, through passage of the Birds and Bees Protection Act (S1856 Hoylman-Sigal). Growing evidence indicates neonics pose neurotoxic risks to wildlife and to people; moreover, New Yorkers' exposure to neonics is widespread and increasing, with potentially untold harms to the health of New York's residents, especially children.

Neonics are neurotoxic insecticides that permanently bind to nerve cellular receptors, causing overstimulation of nerve cells and, eventually, paralysis and death of exposed insects. These same receptors populate the cells of the brain and nervous system across species—including humans, where they are prevalent in sensitive brain areas (such as the cortex, thalamus, and cerebellum) that play a critical role in early growth and development.¹

Exposures to neurotoxic chemicals like neonics during pregnancy and early in life raise special concerns—similar to those raised by lead and mercury—because of the exquisite sensitivity of the developing brain and nervous system. Tragically, experience demonstrates that even miniscule exposures to lead and mercury can lead to lifelong neurological harm, including reduced intelligence (i.e., lower IQ scores), shorter attention spans, and behavioral disruptions.

These concerns are heightened by evidence that neonic exposure in New York is rampant and growing. In 2016, Centers for Disease Control and Prevention (CDC) monitoring found over half the U.S. population had traces of neonics in their urine.² However, a more recent academic study—from 2017 and 2020, testing 171 pregnant women across New York, four additional states, and Puerto Rico—detected neonics *in over 90% of participants*, with higher rates of exposure in Hispanic women.³ Neonic levels were higher than in earlier CDC monitoring, and also *steadily increased over the four-year study* in both level and frequency of detection. This indicates a disturbing trend (and one also observed in wild deer⁴).

Since neonics pass from mother to the developing fetus,⁵ and from breast milk to nursing newborns,⁶ neonic exposures in pregnant and nursing mothers is a significant concern for our children. While further

research will no doubt delineate more fully the health effects of neonics on people, there is already enough concerning evidence to warrant taking immediate action to reduce neonic exposures:

- **Birth defects of the heart** - A study of infants born to mothers in California's San Joaquin valley reported a statistically significant association between living close to agriculture areas that use imidacloprid and having a baby born with Tetralogy of Fallot, a rare heart defect.⁷
- **Birth defects of the brain** - A study of the same population (in California's San Joaquin valley) reported a suggestive association between living near agriculture areas that use imidacloprid and an almost 3-fold elevated risk of having a baby born with anencephaly, lacking parts of the brain and skull.⁸
- **Autism-like symptoms** - A study by NIH-funded researchers from UNC Chapel Hill and UC Davis reported that frequent exposure to imidacloprid during pregnancy was associated with a 2-fold elevated risk of autism spectrum disorder in prenatally exposed children.⁹
- **Acute poisoning reports** - From 2009-2019, the U.S. Environmental Protection Agency (EPA) received more than 1,600 reports of people and pets poisoned with the neonic imidacloprid,¹⁰ with reported symptoms including muscle tremors, difficulty breathing, memory loss, and more—all signs of poisoning with a neurotoxic chemical.¹¹
- **Reduced Sperm Counts and Testosterone Levels** - Studies also link neonic exposure in adults to decreased testosterone levels¹² and significant reductions in sperm count and motility.¹³

Research in animals corroborates harms from neonic exposure across a wide range of mammals. Unpublished testing on neonics—submitted to EPA by pesticide manufacturers for product approvals—reports a wide variety of developmental deficits in prenatally exposed rodents, including statistically significant decreases in the thickness of key areas of the brain (caudate and putamen),¹⁴ altered brain morphology,¹⁵ and significantly delayed and altered reflexes.¹⁶ A recent study of white-tailed deer linked neonic exposures in the womb with decreased body and organ weight, decreased jawbone length, and higher mortality rates for fawns.¹⁷ Research also links neonic exposures in mammals to decreased sperm count, sperm motility, live sperm, and testosterone levels, as well as higher incidence of hypothyroidism and lethargy, delayed reproductive development, and sperm abnormalities.^{18, 19, 20, 21, 22, 23, 24, 25, 26}

The Birds and Bees Protection Act (S1856) eliminates unnecessary neonic uses that account for 80-90% of the chemicals entering the state's environment yearly—specifically, those which in-depth Cornell University research finds provide no economic benefits to users or are replaceable with safer, effective alternatives.²⁷ The act would significantly reduce direct neonic exposure in treated lawns and parks as well as runoff into New York's water sources, which federal and state testing show are extensively polluted with neonics, especially on Long Island.^{28, 29} Drinking water contamination presents a significant exposure concern, given that conventional water treatment generally fails to remove neonics.^{30,31} Infants may be doubly exposed through contaminated baby food as well as formula reconstituted with contaminated tap water.

New York must not repeat the same mistakes we as a nation made with lead and other older neurotoxic chemicals—delaying action while millions of children were exposed and suffered permanent damage. Allowing needless and replaceable neonic uses is simply unjustifiable. Accordingly, we strongly urge passage of the Birds and Bees Protection Act or other decisive government action to reduce New Yorkers' exposure to neurotoxic neonicotinoid pesticides.

Respectfully,

Alycia Halladay, PhD
Chief Science Officer
Autism Science Foundation

Philip J. Landrigan, MD, MSc, FAAP
Director, Program for Global Public Health and the Common Good
Director, Global Observatory on Pollution and Health
Professor of Biology, Schiller Institute for Integrated Science and Society
Boston College, MA
Professor Emeritus of Pediatrics and Preventive Medicine
Icahn School of Medicine at Mount Sinai

Kathleen Nolan, MD, MSL
President, Physicians for Social Responsibility, New York
Co-founder, Concerned Health Professionals of New York
Senior Research Director, Catskill Mountainkeeper

Dr. Stephen de Waal Malefyt, MD
Pediatrician and Director, Albany Children's Environmental Health Center
Albany Medical Center

The following supporters are listed in alphabetical order. For all signatories, academic/institutional affiliation is provided for identification purposes only and does not constitute institutional endorsement unless otherwise indicated.

Natasha Anushri Anandaraja, MD, MPH

Tanya Azarani, MD

Michelle Bamberger, MS, DVM
Retired

David Carpenter, MD
Professor; Director, Institute for Health and the Environment
State University of New York at Albany

Susan Elan, MPH
Department of Public Health
New York Medical College

Sarah Evans, PhD, MPH
Assistant Professor
Icahn School of Medicine at Mount Sinai

Maida Galvez, MD, MPH
Professor
Icahn School of Medicine at Mount Sinai

Julie B. Herbstman, PhD ScM
Professor, Department of Environmental Health Sciences
Director, Columbia Center for Children's Environmental Health
Columbia University Mailman School of Public Health

William Jordan, MD, MPH
Clinical Assistant Professor
Albert Einstein College of Medicine

Carol Kessler, MD, MDiv

Betty Kolod, MD

Bruce P. Lanphear, MD, MPH
Professor of Children's Environmental Health
BC Children's Hospital
Simon Fraser University

Fred Lado, MD, PhD
Director, Epilepsy Division
Northwell Health

Laurie Laugeman, BSN, RN, HNB-BC, HWNC-BC
Sisters of Charity Hospital, Buffalo
Chapter Leader, Western New York Chapter of the American Holistic Nurses Association

Maureen Miller, MD, MPH

Kit O'Malley, MA, RN, HNBC-BC, HWNC-BC, CHT

Robert Oswald, PhD
James Law Professor of Molecular Medicine
Department of Molecular Medicine, Cornell University College of Veterinary Medicine

Micah Resnick, MD
Former Assistant Professor of Pediatrics
Icahn School of Medicine at Mount Sinai
Pediatrician, TriHealth Madeira Pediatrics

Barton Schoenfeld, MD, FACC
Albany Medical Center

Perry Sheffield, MD, MPH
Icahn School of Medicine at Mount Sinai

Peter Sherman, MD, MPH
Chair of Pediatrics
BronxCare Health System

Deborah Socolar, MPH

Rebecca Stern, Ph.D
Postdoctoral Fellow
Harvard T.H. Chan School of Public Health

Adrienne Wald, EdD, MBA RN
Associate Professor of Nursing
Mercy College - Dobbs Ferry, NY

Elie Ward, MSW
Director of Policy & Advocacy
On behalf of:
NYS American Academy of Pediatrics, Chapters 1, 2 & 3

Blair Wylie, MD, MPH
Professor of Obstetrics and Gynecology
Founding Director, The Collaborative for Women's Environmental Health
Columbia University

References:

- ¹ Houchat J-N, Cartereau A, Le Mauff A, Taillebois E, Thany SH. An Overview on the Effect of Neonicotinoid Insecticides on Mammalian Cholinergic Functions through the Activation of Neuronal Nicotinic Acetylcholine Receptors. *International Journal of Environmental Research and Public Health*. 2020; 17(9):3222. <https://doi.org/10.3390/ijerph17093222>.
- ² Ospina M, Wong LY, Baker SE, Serafim AB, Morales-Agudelo P, Calafat AM. Exposure to Neonicotinoid Insecticides in the U.S. General Population: Data from the 2015-2016 National Health and Nutrition Examination Survey. *Environ Res*. 2019 Sep;176:108555. <https://doi.org/10.1016/j.envres.2019.108555>.
- ³ Buckley JP, Kuiper JR, Bennett DH, et al. Exposure to Contemporary and Emerging Chemicals in Commerce among Pregnant Women in the United States: The Environmental influences on Child Health Outcome (ECHO) Program. *Environ Sci Technol*. 2022;56(10):6560-6573. <https://doi.org/10.1021/acs.est.1c08942>.
- ⁴ Dan Gunderson, Data Show Increasing Insecticide Levels in Minnesota Deer, MPR News (Aug. 23, 2022), <https://bit.ly/3RYEcCR>.
- ⁵ Ichikawa G, Kuribayashi R, Ikenaka Y, Ichise T, Nakayama SMM, et al. (2019) LC-ESI/MS/MS analysis of neonicotinoids in urine of very low birth weight infants at birth. *PLOS ONE* 14(7): e0219208. <https://doi.org/10.1371/journal.pone.0219208>.
- ⁶ Zhang D, Lu S. Human Exposure to Neonicotinoids and the Associated Health Risks: A Review. *Environ Int*. 2022;163:107201. <https://doi.org/10.1016/j.envint.2022.107201>.
- ⁷ Carmichael SL, Yang W, Roberts E, Kegley SE, Padula AM, English PB, Lammer EJ, Shaw GM. Residential agricultural pesticide exposures and risk of selected congenital heart defects among offspring in the San Joaquin Valley of California. *Environ Res*. 2014 Nov; 135:133-8. <https://doi.org/10.1016/j.envres.2014.08.030>.
- ⁸ Yang W, Carmichael SL, Roberts EM, Kegley SE, Padula AM, English PB, Shaw GM. Residential agricultural pesticide exposures and risk of neural tube defects and orofacial clefts among offspring in the San Joaquin Valley of California. *Am J Epidemiol*. 2014 Mar 15;179(6):740-8. <https://doi.org/10.1093/aje/kwt324>.
- ⁹ Keil AP, Daniels JL, Hertz-Picciotto I. Autism Spectrum Disorder, Flea and Tick Medication, and Adjustments for Exposure Misclassification: the CHARGE (Childhood Autism Risks from Genetics and Environment) case-control study. *Environ Health*. 2014 Jan 23;13(1):3. <https://doi.org/10.1186/1476-069X-13-3>.
- ¹⁰ Data records obtained via Freedom of Information Act Request, (under "Released Records," select "Specified Incidents for PC 129099 from 1-1-2009 to 4-4-2019"). <https://foiaonline.gov/foiaonline/action/public/submissionDetails?trackingNumber=EPA-HQ-2019-004044&type=request>.

-
- ¹¹ U.S. EPA, Recognition and Management of Pesticide Poisonings: Sixth Edition: 2013, p. 91. https://www.epa.gov/sites/default/files/documents/rmpp_6thed_ch9_otherinsecticides.pdf.
- ¹² Mendy A, Pinney SM. Exposure to Neonicotinoids and Serum Testosterone in Men, Women, and Children. *Environ Toxicol*. 2022;37(6):1521-1528. <https://doi.org/10.1002/tox.23503>.
- ¹³ Hafez EM, Issa SY, Al-Mazroua MK, Ibrahim KT, Rahman SMA (2016) The Neonicotinoid Insecticide Imidacloprid: A Male Reproductive System Toxicity Inducer-Human and Experimental Study. *Toxicol open access* 1:109. <https://bit.ly/3DdBOCP>.
- ¹⁴ EPA 2002. <https://archive.epa.gov/pesticides/chemicalsearch/chemical/foia/web/pdf/129099/129099-0000-00-00a.pdf>.
- ¹⁵ EPA 2007. (MRID Nos. 46028201, 46028202, 47034201). see Table 15, https://www3.epa.gov/pesticides/chem_search/cleared_reviews/csr_PC-060109_9-Mar-07_a.pdf.
- ¹⁶ EPA, HIARC Meeting on Acetamiprid, Briefing Package (Sept. 20, 2001). https://www3.epa.gov/pesticides/chem_search/cleared_reviews/csr_PC-099050_20-Sep-01_a.pdf.
- ¹⁷ Berheim, E.H., Jenks, J.A., Lundgren, J.G. *et al.* Effects of Neonicotinoid Insecticides on Physiology and Reproductive Characteristics of Captive Female and Fawn White-tailed Deer. *Sci Rep* 9, 4534 (2019). <https://doi.org/10.1038/s41598-019-40994-9>.
- ¹⁸ Tariba Lovaković B, Kašuba V, Sekovanić A, Orct T, Jančec A, Pizent A. Effects of Sub-Chronic Exposure to Imidacloprid on Reproductive Organs of Adult Male Rats: Antioxidant State, DNA Damage, and Levels of Essential Elements. *Antioxidants (Basel)*. 2021;10(12):1965. Published 2021 Dec 8. <https://doi.org/10.3390%2F antioxidants10121965>.
- ¹⁹ Lonare M, Kumar M, Raut S, et al. Evaluation of Ameliorative Effect of Curcumin on Imidacloprid-Induced Male Reproductive Toxicity In Wistar Rats. *Environ Toxicol*. 2016;31(10):1250-1263. <https://doi.org/10.1002/tox.22132>.
- ²⁰ Hafez et al. 2016.
- ²¹ Bagri P, Kumar V, Sikka AK. An in Vivo Assay of the Mutagenic Potential of Imidacloprid Using Sperm Head Abnormality Test and Dominant Lethal Test. *Drug Chem Toxicol*. 2015;38(3):342-348. <https://doi.org/10.3109/01480545.2014.966832>.
- ²² Bal R, Naziroğlu M, Türk G, et al. Insecticide Imidacloprid Induces Morphological and DNA Damage Through Oxidative Toxicity on the Reproductive Organs of Developing Male Rats. *Cell Biochem Funct*. 2012;30(6):492-499. <https://doi.org/10.1002/cbf.2826>.
- ²³ Bal R, Türk G, Tuzcu M, et al. Assessment of Imidacloprid Toxicity on Reproductive Organ System of Adult Male Rats. *J Environ Sci Health B*. 2012;47(5):434-444. <https://doi.org/10.1080/03601234.2012.663311>.
- ²⁴ Berheim et al. 2019.
- ²⁵ Najafi G, Sc D, Razi M, Hoshyar A, Shahmohamadloo S, Feyzi S. The Effect of Chronic Exposure with Imidacloprid Insecticide on Fertility in Mature Male Rats. *Int J FertilSteril*. 2009. 9. <https://bit.ly/3yGr0vQ>.
- ²⁶ U.S. Environmental Protection Agency, Data Evaluation Report (DER) for Thiamethoxam: Developmental Neurotoxicity Study, <https://bit.ly/3TlXKSL>; MRID 47034201, 46028202.
- ²⁷ Grout T.A., Koenig P.A., Kapuvari J.K. Neonicotinoid Insecticides in New York State: Economic Benefits and Risk to Pollinators. Jun 23, 2020. <https://bit.ly/3esfOMp>.
- ²⁸ Importantly, nearly all federal and state testing is for imidacloprid alone, not the other four major neonics in use in New York State. Since imidacloprid is estimated to be only the second-most used neonic chemical in the state, actual neonic water contamination is likely much higher than reported. Pierre Mineau, Impacts of Neonics in New York Water: Their Use and Threats to the State's Aquatic Ecosystems (Sep. 2019), <https://on.nrdc.org/2lXs000>.
- ²⁹ Fisher IJ, Phillips PJ, Bayraktar BN, Chen S, McCarthy BA, Sandstrom MW. Pesticides and Their Degradates in Groundwater Reflect Past Use and Current Management Strategies, Long Island, New York, USA. *Sci Total Environ*. 2021;752:141895. <https://doi.org/10.1016/j.scitotenv.2020.141895>.
- ³⁰ Klarich K.L., Pflug N.C., DeWald E.M., Hladik M.L., Kolpin D.W., Cwiertny D.M., LeFevre G.H. Occurrence of Neonicotinoid Insecticides in Finished Drinking Water and Fate During Drinking Water Treatment. *Environ. Sci. Technol. Lett*. 2017 4, 5, 168-73. <https://doi.org/10.1021/acs.estlett.7b00081>.
- ³¹ See, e.g., Suffolk County Water Authority, 2022 Drinking Water Quality Report, http://s1091480.instanturl.net/dwqr2022/AWQR_2022_FINAL.pdf (showing the neonics imidacloprid appears in tap water in some of Suffolk County's largest water distribution areas).